

Author Index

Abernathy JR. see Perova NV Aiken J. see Lazarus R Albert MS. How does education affect cognitive function? (editorial), 76-78 Aldoori WH, Giovannucci EL, Rimm EB, Wing AL, Trichopoulos DV, Willett WC. A prospective study of alcohol, smoking, caffeine, and the risk of symptomatic diverticular disease in men, 221-228 Allender PS. see Hollis JF Anderson CV. see Clark K Appel LJ Espeland M, Whelton PK, Dolecek T, Kumanyika S, Applegate WB, Ettinger WH Jr, Kostis JB, Wilson AC, Lacy C, Miller ST, Trials of Hypertension Prevention (TOHP) Collaborative Research Group. Trial of nonpharmacologic intervention in the elderly (TONE): design and rationale of a blood pressure control trial, 119-129 Hebert PR, Cohen JD, Obarzanek E, Yamamoto M, Buring J, Stevens V, Kirchner K, Borhani NO. Baseline characteristics of participants in phase II of the Trials of Hypertension Prevention (TOHP II), 149-155

see Kahn HA see Sacks FM

Applegate WB see Appel LJ

see Yamamoto ME

Armstrong BG. Comparing standardized mortality ratios, 60–64

Arnold JE, Rohan T, Howe G, LeBlanc M. Reproducibility and validity of a food-frequency questionnaire designed for use in girls age 7 to 12 years, 369–377

Atherosclerosis Risk in Communities (ARIC) Study Investigators see Duncan BB see Folsom AR Augier A. see Clavel-Chapelon F

B Barbone F. see Franceschi S Barnes RW. see Folsom AR Barrier P. see Crawford PB Bartko JJ. see Farmer ME Batey DM. see Lasser VI Beard CM, Spencer RJ, Weiland LH, O'Fallon WM,
Melton LJ III. Trends in colorectal cancer over a half
century in Rochester, Minnesota, 1940 to 1989, 210–
214
Belden LK. see Lasser VI
Benichou J. see D'Avanzo B
Bergman RN. see Wagenknecht LE
Bild DE
see Ives DG

see Psaty BM Blair N. see Hiller R Bolt RJ. see Hebert PR Borhani NO

see Appel LJ see Hebert PR see Hollis JF see Whelton PK see Yamamoto ME

Borok GM. see Wagenknecht LE Boyle CA. see Decouflé P Bracken MB. see Greenberg RS Branch LG. see Gaziano JM Brand L. see Layde PM Brown CP. see Matanoski GM Bray GA. see Sacks FM Brémond A. see Clavel-Chapelon F

Brett KM
Madans JH. Long-term survival after coronary heart
disease: comparisons between men and women in a
national sample, 25–32

see Madans JH Brewer AA. see Lasser VI Buffler PA. see Greenberg RS Buhari AM. see Massoudi MS

Burchfiel CM, Curb D, Rodriguez BL, Yano K, Hwang L-J, Fong K-O, Marcus EB. Incidence and predictors of diabetes in Japanese-American men: the Honolulu heat program, 33–43

Buring JE

see Appel LJ

see Gaziano JM

see Manson JE

see Whelton PK

Burke GL. see Psaty BM

Burt VL. see Dalager NA

Burton TC. see Hiller R

Busby MG. see Cooper GS Byrne CD. see Wild SH

C

Caan BJ. see Slattery ML Califf RM. see Peterson ED

Cao G. see Hauck FR

Carlson W. see Hanrahan JP

Carpenter MA. see Duncan BB

Chambless LE. see Duncan BB

Charleston J. see Hollis JF

Chen Y-X. see Zhang Z-F

Choo PW. see Hanrahan JP

Cirillo M. see Trevisan M

Clark K, Sowers MF, Wallace RB, Jannausch ML, Lemke J, Anderson CV. Age-related hearing loss and bone mass in a population of rural women aged 60 to 85 years, 8-14

Clavel-Chapelon F, Launoy G, Auqier A, Gairard B, Brémond A, Piana L, Lansac J, Renaud R. Reproductive factors and breast cancer risks: effect of age at diagnosis, 315–320

Cleghorn FR, Maybank KA, Jack N, Pate E, Mingle J, Levine PH, Manns A. Comparison of HHV-6 antibody titers in West Africa and the Carribean, 497–500

Cohen JD

see Appel LJ

see Hebert PR

see Whelton PK

see Yamamoto ME

Colditz GA. see Gaziano JM

Cook NR

see Hebert PR

see Manson JE

see Whelton PK

Cooper GS, Busby MG, Fairchild AP. Measurement of lactose consumption. Reliability and comparison of two methods, 473–477

Coory M. An investigation into the disparity between Australian Aboriginal and Caucasian perinatal mortality rates, 393–399

Corey PN. see Sass-Kortsak AM

Costantino JP. see Massoudi MS

Cramer DW, Xu H. Epidemiologic evidence of uterine growth factors in the pathogenesis of ovarian cancer, 310–314

Crawford PB, Obarzanek E, Schreiber GB, Barrier P, Goldman S, Frederick M, Sabry ZI. The effects of race, household income, and parental education on nutrient intakes of 9- and 10-year-old girls: NHLBI growth and health study, 360–368

Crouse JR III. see Duncan BB Crowley PM. see Ives DG Cruise G. see Ives DG

Curb D. see Burchfiel CM

Cutler JA

Progress in life-style intervention for prevention and treatment of high blood pressure (editorial), 165–167

see Hebert PR

see Whelton PK

D

Dalager NA, Kang HK, Burt VL, Weatherbee L. Hodgkin's disease and Vietnam service, 400-406

Dalcin A. see Lasser VI

Daltroy LH. see Karlson EW

Danielson E. see Lasser VI

Darlington GA. see Jaglal SB

DASH Investigators. see Sacks FM

D'Avanzo B, La Vecchia C, Negri E, DeCarli A, Benichou J. Attributable risks for bladder cancer in Northern Italy, 427-431

Davis CE. see Perova NV

Decarli A

see D'Avanzo B

see Franceschi S

Decouflé P, Boyle CA. The relationship between maternal education and mental retardation in 10-year-old children, 347-353

Deev AD. see Perova NV

Dersimonian R. see Strickler H

Dolecek T. see Appel LJ

Doxiadis S. see Petridou E

Duncan BB, Chambless LE, Schmidt MI, Szklo M, Folsom AR, Carpenter MA, Crouse JR III, Atherosclerosis Risk in Communities (ARIC) Study Investigators. Correlates of body fat distribution: variation across categories of race, sex, and body mass in the atherosclerosis risk in communities study, 192–200

Dyer A. see Trevisan M

E

Eckfeldt JH. see Folsom AR

Edwards SL. see Slattery ML Espeland M. see Appel LJ

Ettinger WH Jr. see Appel LJ

Evans MA

see Lasser VI

see Sacks FM

Eye Disease Case-Control Study Group. see Hiller R

F

Faich GA. see Hanrahan JP

Fairchild AP. see Cooper GS

Farber MD. see Hiller R

Farmer ME, Kittner SJ, Rae DS, Bartko JJ, Regier DA. Education and change in cognitive function: the epidemiological catchment area study, 1–7

Ferguson JA. see Matanoski GM

Ferraroni M. see Franceschi S

Filiberti R. see Franceschi S

Fitzpatrick AL. see Ives DG

Flytzani V. see Petridou E

Foley TP Jr. see Massoudi MS

Folsom AR

Ma J, Eckfeldt JH, Nieto FJ, Metcalf PA, Barnes RW, Atherosclerosis Risk in Communities (ARIC) Study Investigators. Low serum albumin: association with diabetes mellitus and other cardiovascular risk factors but not with prevalent cardiovascular disease or carotid artery intima-media thickness, 186–191 see Duncan BB

Fong K-O. see Burchfiel CM

Fontham E. see Matanoski GM

Fortmann SP. see Wild SH

Fouad M. see Hollis JF

Franceschi S

Barbone F, Negri E, Decarli A, Ferraroni M, Filiberti R, Giacosa A, Gnagnarella P, Nanni O, Salvini S, La Vecchia C. Reproducibility of an Italian food frequency questionnaire for cancer studies: results for specific nutrients, 69–75

Franceschi S. see La Vecchia C

Frederick M. see Crawford PB

G

Gairard B. see Clavel-Chapelon F

Gallagher K. see Payne SMC

Garcia G. see McPherson RS

Gaziano JM. see Manson JE

Gaziano JM, Manson JE, Branch LG, Colditz GA, Willett WC, Buring JE. A prospective study of consumption of carotenoids in fruits and vegetables and decreased cardiovascular mortality in the elderly, 255–260

Gazmararian JA, James SA, Lepkowski JM. Depression in black and white women. The role of marriage and socioeconomic status, 455–463

Giacosa A. see Franceschi S

Giovannucci EL. see Aldoori WH

Gliksman M. see Lazarus R

Gnagnarella P. see Franceschi S

Goff DC Jr, Nichaman MZ, Ramsey DJ, Meyer PS, Labarthe DR. A population-based assessment of the use and effectiveness of thrombolytic therapy: the Corpus Christi Heart Project, 171-178 Goldman S. see Crawford PB

Goodwin Sousoulas B. see Hollis JF

Gragoudas ES. see Hiller R

Graham S. see Zhang Z-F

Greenberg RS, Buffler PA, Hinman AR, Swanson GM, Matanoski GM, Nasca PC, Bracken MB, Schoenbach VJ. Commentary—American College of Epidemiology statement of principles (ACE section), 503–504

Greineder D. see Hanrahan IP

Gubbio Collaborative Study Group. see Trevisan M

Gunter EW. see Hiller R

H

Haffner S. see Wagenknecht LE

Haller J. see Hiller R

Hanis CL. see McPherson RS

Hanrahan JP, Choo PW, Carlson W, Greineder D, Faich GA, Platt R. Terfenadine-associated ventricular arrhythmias and QT_c interval prolongation: a retrospective cohort comparison with other antihistamines among members of a health maintenance organization, 201–209

Hauck FR, White L, Cao G, Woolf N, Strauss K. Inaccuracy of self-reported weights and heights among American Indian adolescents, 386–392

Hebert PR

Bolt RJ, Borhani NO, Cook NR, Cohen JD, Cutler JA, Hollis JF, Kuller LH, Lasser NL, Oberman A, Miller ST, Morris C, Whelton PK, Hennekens CH, Trials of Hypertension Prevention (TOHP) Collaborative Research Group. Design of a multicenter trial to evaluate long-term life-style intervention in adults with high-normal blood pressure levels: trials of hypertension prevention (phase II), 130–139

see Appel LJ

see Kahn HA

Hennekens CH

see Hebert PR

see Manson JE

see O'Donnell CJ

see Yamamoto ME

Henkin L. see Wagenknecht LE

Hertz T. see Payne SMC

Hiller R, Seigel D, Sperduto RD, Blair N, Burton TC, Farber MD, Gragoudas ES, Gunter EW, Haller J, Seddon JM, Sowell AL, Yannuzzi LA, Eye Disease Case-Control Study Group. Serum zinc and serum lipid profiles in 778 adults, 490–496

Hinman AR. see Greenberg RS

Hirlinger M. see Hollis JF

Hollis JF

Satterfield S, Smith F, Fouad M, Allender PS, Borhani N, Charleston J, Hirlinger M, King N, Schultz R, Goodwin Sousoulas B, Trials of Hypertension Prevention (TOHP) Collaborative Research Group. Recruitment for phase II of the trials of hypertension prevention: effective strategies and predictors of randomization, 140–148

see Hebert PR Honjo S. see Todoroki I Hoover DR. see Strickler H Howard G. see Wagenknecht LE Howe G. see Arnold JE

Hwang L-J. see Burchfiel CM

I

Imanishi K. see Todoroki I
Irving SH. see Perova NV
Ives DG, Fitzpatrick AL, Bild DE, Psaty BM, Kuller LH,
Crowley PM, Cruise G, Theroux S. Surveillance and
ascertainment of cardiovascular events: the
Cardiovascular Health Study, 278–285

Jack N. see Cleghorn FR
Jacobs DR Jr. see Slattery M
Jacques PF, Sulsky SI, Perrone GA, Jenner J, Schaefer EJ.
Effect of vitamin C supplementation on lipoprotein cholesterol, apolipoprotein, and triglyceride concentrations, 52–59
Jaglal SB, Kreiger N, Darlington GA. Lifetime occupational physical activity and risk of hip fracture in women, 321–324
James SA

see Gazmararian JA
see Matanoski GM
Jannausch ML. see Clark K
Jenkins B. see Matanoski GM
Jenner J. see Jacques PF

K

Kahn HA, Whelton PK, Appel LJ, Kumanyika SK, Meneses JL, Hebert PR, Woods M. Validity of 24-hour dietary recall interviews conducted among volunteers in an adult working community, 484–489 Kanellopoulou G. see Petridou E

Kang HK see Dalager NA see Watanabe KK Karanja N. see Sacks FM

Karlson EW, Sanchez-Guerrero J, Wright EA, Lew RA, Daltroy LH, Katz JN, Liang MH. A connective tissue disease screening questionnaire for population studies, 297–302

297-302
Katsurada M. see Todoroki I
Katz JN. see Karlson EW
Kerber RA. see Slattery ML
Kiley JE. see Whelton PK
King N. see Hollis JF
Kirchner KA
see Appel LJ
see Yamamoto ME

Kittner SJ see Farmer ME see Psaty BM

Klag MJ. see Yamamoto ME Kohl HW III. see McPherson RS Kono S. see Todoroki I Kostis JB. see Appel LJ

Kreiger N. see Jaglal SB Krogh V. see Trevisan M

Krogh V. see Trevisan
Kuller LH
see Hebert PR
see Ives DG
see Massoudi MS
see Psaty BM
see Whelton PK

see Whelton PK
Kumanyika S
see Appel LJ
see Kahn HA
see Lasser VI
see Matanoski GM

L
Labarthe DR. see Goff DC Jr
Lacy C. see Appel LJ
Lakatos E. see Yamamoto ME
Lancashire R. see Sorahan T
Lansac J. see Clavel-Chapelon F
Lasser NL. see Hebert PR

Lasser VI, Raczynski JM, Stevens VJ, Mattfeldt-Beman MK, Kumanyika S, Evans M, Danielson E, Dalcin A, Batey DM, Belden LK, Brewer AA, Trials of Hypertension Prevention (TOHP) Collaborative Research Group. Trials of hypertension prevention, phase II: structure and content of the weight loss and dietary sodium reduction interventions, 156–164

Launoy G. see Clavel-Chapelon F Laurenzi M. see Trevisan M

La Vecchia C

Levi F, Lucchini F, Franceschi S. Trends in mortality from nonneoplastic gallbladder disease, 215–220 see D'Avanzo B see Franceschi S see Tavani A Laws A. see Wild SH

Layde PM, Nordstrom DL, Stueland D, Brand L, Olson KA. Machine related occupational injuries in farm residents, 419–426

Lazarus R, Wilson A, Gliksman M, Aiken J. Repeatability of nutrient intakes estimated by a semiquantitative food frequency questionnaire in elderly subjects, 65–68

LeBlanc M. see Arnold JE

Lemke J. see Clark K

Lepkowski JM. see Gazmararian JA

Levi F. see La Vecchia C

Levine PH. see Cleghorn FR

Lew RA. see Karlson EW

Liang MH. see Karlson EW

Liao C-S. see Zhang Z-F

Lin P-H. see Sacks FM

Linn S. The injury severity score—importances and uses, 440–446

Losina H. see Payne SMC

Lucchini F. see La Vecchia C

M

Ma J. see Folsom AR

Malamou H. see Petridou E

Manns A. see Cleghorn FR

Manson JE

Gaziano JM, Spelsberg A, Ridker PM, Cook NR, Buring JE, Willett WC, Hennekens CH, WACS Research Group. A secondary prevention trial of antioxidant vitamins and cardiovascular disease in women: rationale, design, and methods, 261–269 see Gaziano JM

Marcus EB. see Burchfiel CM

Marcus PM, Newcomb PA, Young T, Storer BE. The association of reproductive and menstrual characteristics and colon and rectal cancer risk in Wisconsin women, 303–309

Marshall J. see Zhang Z-F

Massoudi MS, Meilahn EN, Orchard TJ, Foley TP Jr, Kuller LH, Costantino JP, Buhari AM. Prevalence of thyroid antibodies among healthy middle-aged women: findings from the thyroid study in healthy women, 229–233

Matanoski GM

Nasca PC, Swanson GM, Vernon SW, Fontham E, Brown CP, Ferguson JA, James SA, Jenkins B, Mays VM, Nwangwu JT, Reynolds GH, Kumanyika SK, Schoenbach VJ, Suarez L. Statement of principles: Epidemiology and minority populations (ACE section), 505–508

see Greenberg RS

Mattfeldt-Beman MK. see Lasser VI

Maybank KA. see Cleghorn FR

Mayer EJ. see Wagenknecht LE

Mays VM. see Matanoski GM

McCullough M. see Sacks FM

McPherson RS, Kohl HW III, Garcia G, Nichaman MZ, Hanis CL. Food-frequency questionnaire vaqlidation among Mexican-Americans: Starr County, Texas, 378– 385

Meilahn EN. see Massoudi MS

Melton LJ III. see Beard CM

Meneses JL. see Kahn HA

Metcalf PA. see Folsom AR

Meyer PS. see Goff DC Jr

Miller ST

see Appel LJ

see Hebert PR

Mingle J. see Cleghorn FR

Mittelmark M. see Psaty BM

Moore TJ. see Sacks FM

Morris C. see Hebert PR

Murphy SP. Food frequency questionnaires: how far have we come and where are we going? (editorial), 413

N

Nanni O. see Franceschi S

Nasca PC

see Greenberg RS

see Matanoski GM

Negri E

see D'Avanzo B

see Franceschi S

see Tavani A

Neutel CI. Risk of traffic accident injury after a prescription

for a benzodiazepine, 239-244

Newcomb PA. see Marcus PM

Nichaman MZ

see Goff DC Jr

see McPherson RS

Nieto FJ. see Folsom AR

Nishikawa H. see Todoroki I

Nordstrom DL. see Layde PM

Nwangwu JT. see Matanoski GM

0

Obarzanek E

see Appel LJ

see Crawford PB

see Sacks FM

Oberman A

Changing of the guard (editorial), 82

see Hebert PR

Oddleifson S. see Payne SMC

O'Donnell CJ, Ridker PM, Hennekens CH. Thrombolytic therapy for acute myocardial infarction: applying lessons from randomized trials to community practice (editorial), 250–252

O'Fallon WM. see Beard CM Oganov RG. see Perova NV

Ogawa S. see Todoroki I

Olson KA. see Layde PM

Omenn GS. What accounts for the association of vegetables and fruits with lower incidence of cancers and coronary heart disease? (editorial), 333–335

Orchard TJ. see Massoudi MS

P

Pantelakis S. see Petridou E

Payne SMC, Seage GR III, Oddleifson S, Gallagher K, vanBeuzekom M, Losina H, Hertz T. Using administratively collected hospital discharge data for AIDS surveillance, 337–346

Pate E. see Cleghorn FR Peck I. see Sorahan T

Perova NV, Oganov RG, Williams DH, Irving SH,
Abernathy JR, Deev AD, Shestov DB, Zhukovsky GS,
Davis CE, Tyroler HA. Association of high-densitylipoprotein cholesterol with mortality and other risk
factors for major chronic noncommunicable diseases in
samples of US and Russian men, 179–185

Perrone GA. see Jacques PF

Peterson ED, Califf RM. The complexity of studying coronary heart disease in women (editorial), 79-81

Petridou E, Malamou H, Doxiadis S, Pantelakis S, Kanellopoulou G, Toupadaki N, Trichopoulou A, Flytzani V, Trichopoulos D. Blood lipids in Greek adolescents and their relation to diet, obesity, and socioeconomic factors, 286–291

Piana L. see Clavel-Chapelon F

Platt R. see Hanrahan IP

Potter JD. see Slattery ML

Price TR. see Psaty BM

Prior P. see Sorahan T

Proschan MA. see Sacks FM

Psaty BM

Kuller LH, Bild D, Burke GL, Kittner SJ, Mittelmark M, Price TR, Rautaharju PM, Robbins J. Methods of assessing prevalent cardiovascular disease in the cardiovascular health study, 270–277 see Ives DG

R

Raczynski JM. see Lasser VI Rae DS. see Farmer ME Ramsey DJ. see Goff DC Jr Rautaharju PM. see Psaty BM
Regier DA. see Farmer ME
Renaud R. see Clavel-Chapelon F
Rewers M. see Wagenknecht LE
Reynolds GH. see Matanoski GM
Ridker PM
see Manson JE
see O'Donnell CJ

Rimm EB. see Aldoori WH Robbins J. see Psaty BM Robertson JM. see Sass-Kortsak AM

Rodriguez BL. see Burchfiel CM

Rohan T. see Arnold JE

Roseman J. Epidemiology: the logic of modern medicine (book review), 501

S

Saad MF. see Wagenknecht LE Sabry ZI. see Crawford PB Sacks FM

Dash Investigators. Letter, 502

Obarzanek E, Windhauser MM, Svetkey LP, Vollmer WM, McCullough M, Karanja N, Lin P-H, Steele P, Proschan MA, Evans MA, Appel LJ, Bray GA, Vogt TM, Moore TJ, DASH Investigators. Rationale and design of the Dietary Approaches to Stop Hypertension trial (DASH): a multicenter controlled-feeding study of dietary patterns to lower blood pressure, 108–118

see Whelton PK see Yamamoto ME

Sakurai Y. see Todoroki I

Salvini S. see Franceschi S

Sanchez-Guerrero J. see Karlson EW

Sass-Kortsak AM, Corey PN, Robertson JM. An investigation of the association between exposure to styrene and hearing loss, 15–24

Satterfield S

see Hollis JF

see Whelton PK

Savage PJ. see Wagenknecht LE

Schaefer EJ. see Jacques PF

Schmidt MI. see Duncan BB

Schoenbach VJ

see Greenberg RS

see Matanoski GM

Schreiber GB. see Crawford PB

Schultz R. see Hollis JF

Seage GR III. see Payne SMC

Seddon JM. see Hiller R

Shestov DB. see Perova NV

Seigel D. see Hiller R

Selby J. see Wagenknecht LE

517

Shinchi K. see Todoroki I

Slattery M, Jacobs DR Jr. Assessment of ability to recall physical activity of several years ago, 292-296

Slattery ML, Edwards SL, Caan BJ, Kerber RA, Potter JD. Response rates among control subjects in casecontrol studies, 245-249

Smith F. see Hollis JF

Sorahan T, Lancashire R, Prior P, Peck I, Stewart A. Childhood cancer and parental use of alcohol and tobacco, 354-359

Sowell AL. see Hiller R

Sowers MF. see Clark K

Spelsberg A. see Manson JE

Spencer RJ. see Beard CM

Sperduto RD. see Hiller R

Stamler J. see Trevisan M

Steele P. see Sacks FM

Stevens V. see Appel LJ

Stevens VJ. see Lasser VI

Stewart A. see Sorahan T

Storer BE. see Marcus PM

Strauss K. see Hauck FR

Strickler H, Hoover DR, Dersimonian R. Problems in intepreting HIV sentinel seroprevalence studies, 447-454

Stueland D. see Layde PM

Suarez L. see Matanoski GM

Sulsky SI. see Jacques PF

Sun M. see Zhang Z-F

Svetkey LP. see Sacks FM

Swanson GM

see Greenberg RS

see Matanoski GM

Szklo M. see Duncan BB

Tang S-L. see Zhang Z-F

Tavani A, Negri E, La Vecchia C. Selected diseases and risk of cataract in women: a case-control study from Northern Italy, 234-238

Taylor JO

see Whelton PK

see Yamamoto ME

Theroux S. see Ives DG

Todoroki I, Kono S, Shinchi K, Honjo S, Sakurai Y, Wakabayashi K, Imanishi K, Nishikawa H, Ogawa S, Katsurada M. Relationship of cigarette smoking, alcocol use, and dietary habits with sigmoid colon adenomas, 478-483

Toupadaki N. see Petridou E

Trevisan M, Krogh V, Cirillo M, Laurenzi M, Dyer A. Stamler J, Gubbio Collaborative Study Group. Red blood sodium and potassium concentration and blood pressure: the Gubbio population study, 44-51

Trials of Hypertension Prevention (TOHP) Collaborative

Research Group

see Appel LJ

see Hebert PR

see Hollis JF

see Lasser VI

see Whelton PK

see Yamamoto ME

Trichopoulos DV

see Aldoori WH

see Petridou E

Trichopoulou A. see Petridou E

Tyroler HA. see Perova NV

V

vanBeuzekom M. see Payne SMC Varady AN. see Wild SH Vernon SW. see Matanoski GM Vogt TM. see Sacks FM Vollmer WM. see Sacks FM

W

WACS Research Group. see Manson JE

Wagenknecht LE, Mayer EJ, Rewers M, Haffner S, Selby J, Borok GM, Henkin L, Howard G, Savage PJ, Saad MF, Bergman RN, Hamman R. The Insulin Resistance Atherosclerosis Study (IRAS). Objectives, design, and recruitment results, 464-472

Wakabayashi K. see Todoroki I

Wallace RB. see Clark K

Watanabe KK, Kang HK. Military service in Vietnam and the risk of death from trauma and selected cancers, 407-412

Weatherbee L. see Dalager NA

Weiland LH. see Beard CM

Whelton PK

see Appel LJ

see Hebert PR

see Kahn HA

Whelton PK, Buring J, Borhani NO, Cohen JD, Cook N, Cutler JA, Kiley JE, Kuller LH, Satterfield S, Sacks FM, Taylor JO, Trials of Hypertension Prevention (TOHP) Collaborative Research Group. The effect of potassium supplementation in persons with a highnormal blood pressure: results from phase I of the trials of hypertension prevention (TOHP), 85-95

White L. see Hauck FR

Wild SH, Laws A, Fortmann SP, Varady AN, Byrne CD. Mortality from coronary heart disease and stroke for six ethnic groups in California, 1985 to 1990, 432–439

Willett WC

see Aldoori WH

see Gaziano JM

see Manson JE

Williams DH. see Perova NV

Williams OD. Total mortality and mortality from heart disease, cancer, and stroke from 1950 to 1987 in 27 countries: highlights of trends and their interrelationships among causes of death (book review), 417

Wilson A. see Lazarus R Wilson AC. see Appel LJ Windhauser MM. see Sacks FM Wing AL. see Aldoori WH Woods M. see Kahn HA Woolf N. see Hauck FR Wright EA. see Karlson EW

X Xu H. see Cramer DW Xu J-L. see Zhang Z-F Y

Yamamoto M. see Appel LJ

Yamamoto ME, Applegate WB, Klag MJ, Borhani NO, Cohen JD, Kirchner KA, Lakatos E, Sacks FM, Taylor JO, Hennekens CH, Trials of Hypertension Prevention (TOHP) Collaborative Research Group. Lack of blood pressure effect with calcium and magnesium suplementation in adults with high-normal blood pressure: results from phase I of the trials of hypertension prevention (TOHP), 96–107

Yang X-Z. see Zhang Z-F Yannuzzi LA. see Hiller R Yano K. see Burchfiel CM Young T. see Marcus PM Yu S-Z. see Zhang Z-F

Zielezny M. see Zhang Z-F

Zahm SH. Epidemiologic research on Vietnam veterans: difficulties and lessons learned (editorial), 414–416
Zhang Z-F, Graham S, Yu S-Z, Marshall J, Zielezny M, Chen Y-X, Sun M, Tang S-L, Liao C-S, Xu J-L, Yang X-Z. *Trichomonas vaginalis* and cervical cancer: a prospective study in China, 325–332
Zhukovsky GS. see Perova NV



Subject Index

A Abbreviated Injury Scale (AIS) development and attributes of, 440-446 **Aborinines** Australian perinatal mortality rates in, 393 Injury Severity Score method for retrospective comparison of, 440-446 Acquired immunodeficiency syndrome (AIDS). see also Human immunodeficiency virus sentinel seroprevalence studies estimating HIV prevalence from, 447-454 surveillance administratively collected hospital discharge data for, Adiposity central correlates of, 192-200 Administrative databases of hospital discharge of AIDS patients in AIDS surveillance, 337-346 Adolescent(s) American Indian self-reporting of weights and heights among, 386-392 blood lipids in, 286-291 nutrition and nutrient intakes of 9- and 10-year-old girls, 360-Adult working community validity of 24-hour dietary recall interviews in, 484-489 African-Americans. see also Blacks coronary heart disease and stroke-related mortality for in California, 1985 to 1990, 432-439 and correlates of body fat distribution, 192-200 Age groups at breast cancer diagnosis effect of, 315-320 cognitive function correlation with

and HHV-6 antibody titers in West Africa and the Caribbean, 497-500 and red blood cell sodium and potassium concentration effects on blood pressure, 44-51 and thyroid antibody prevalence in middle-aged women, 229-233 and traffic accident injury risk after prescription for benzodiazepine, 239-244 coronary heart disease and stroke-related mortality for in California, 1985 to 1990, 432-439 Agent Orange as Hodgkin's disease factor, 400-406 Agriculture high rates of occuptaional injury in, 419-426 AIS. see Abbreviated Injury Scale Albumin low serum concentration associated with diabetes mellitus and other cardiovascular risk factors, 186-191 Alcohol. see also Ethanol in diverticular disease in men, 221-228 parental use correlated with childhood cancer, 354-359 and reproducibility of food frequency questionnaire for cancer studies, 69 and sigmoid colon adenomas, 478-483 American College of Epidemiology Statement of principles: epidemiology and minority populations (ACE section), 505-508 commentary, statement of principles, 503-504 American Indian adolescents self-reporting of weights and heights among, 386-392 Anatomic and physiological injury combined scales of Injury Severity Score method and, 440-446 Angina pectoris prevalence assessing in elderly, 270-277 Announcements, 83, 169 Antibody titers HHV-6 in West Africa and the Caribbean 497 Antihistamine-associated ventricular arrhythmias QT_c interval prolongation with, 201–209

(editorial), 1-7

hearing loss related to

8-14

assessed via epidemiologic catchment study

and bone mass in rural women aged 60 to 85 years,

Antioxidants

consumption of

in fruits and vegetables, 255–260, 333–335 (editorial) secondary prevention trial of in women, 261–269

Anxiolytics

prescription for

risk of traffic accident injury after, 239-244

Apolipoproteins

vitamin C supplementation effect on, 52-59

Arrhythmias

terfenadine-associated ventricular QT_c interval prolongation with, 201-209

Ascertainment

of cardiovascular events

in Cardiovascular Health Study, 278-285

Ascorbic acid. see also Vitamin C

supplementation

effect of on lipoprotein cholesterol, apolipoprotein, and triglycerides, 52-59

Asian Indians

coronary heart disease and stroke-related mortality for in California, 1985 to 1990, 432-439

Asian-Americans. see also Japanese Americans incidence and predictors of diabetes in, 33-43

Asymptotic precision

in comparing standardized mortality ratios, 60-64

Atherosclerosis

and correlates of body fat distribution, 192-200 low serum albumin associated with, 186-191

Attributable risks

for bladder cancer in northern Italy 427-431

B

Benzodiazepines

prescription for

risk of traffic accident injury after, 239-244

Beta-carotene

for cardiovascular disease prevention in women, 261-269 consumption of

in fruits and vegetables, 255-260, 333-335 (editorial) in food frequency questionnaire designed for use in girls age 7 to 12 years, 369-377

and repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire 65-68

Biases

comparing standardized mortality ratios, 60-64 time-dependent

in HIV sentinel seroprevalence studies, 447-454

Birth weight

neonatal

and disparity between Australian Aboriginal and Caucasian perinatal mortality rates, 393–399 and maternal consumption of cigarettes, 354–359 Blacks. see also African-Americans

mothers

and maternal education correlated with mental retardation in 10-year-old children, 347–353 and nutrient intakes of 9- and 10-year-old girls, 360–

women

role of marriage and socioeconomic status on depression in, 455–463

Bladder cancer

attributable risks for in northern Italy 427-431

Blood lipids

in Greek adolescents

relation to diet, obesity, and socioeconomic factors, 286-291

Blood pressure. see also Hypertension

high-normal

calcium supplementation in, 96–107 magnesium supplementation in, 96–107 potassium supplementation in, 85–95

progress in life-style intervention for prevention and treatment of (editorial), 165-167

Body fat. see Fat, body

Body height

in American Indian adolescents inaccuracy in self-reporting of, 386-392

Body mass index

and cataract risk in women, 234–238 234 and correlates of body fat distribution, 192–200 in self-reporting of weights and heights among American Indian adolescents, 386–392

Body weight. see also Obesity

in American Indian adolescents inaccuracy in self-reporting of, 386-392

Bone mass

related to age-related hearing loss in elderly rural women, 8-14

Breast cancer

risks

and reproductive factors, 315-320

C

Caffeine

in diverticular disease in men, 221-228

Calcium

and nutrient intakes of 9- and 10-year-old girls, 360-368

supplementation

in persons with high-normal blood pressure, 96-107

Cancer

HDL cholesterol association with in US and Russian men, 179-185

heart disease, and stroke, total mortality and mortality from, from 1950 to 1987 in 27 countries (book review), 417

risk

and association of reproductive and menstrual characteristics, 303-309

studies

reproducibility of Italian food frequency questionnaire for, 69

selective

and Vietnam service, 407-412

Cardiovascular disease

antioxidant vitamins for prevention of, 261-269 low serum albumin associated with, 186-191 mortality in elderly

decreased by carotenoid consumption in fruits and vegetables, 255-260, 333-335 (editorial)

surveillance and ascertainment of, 278-285

Cardiovascular Health Study

on surveillance and ascertainment of cardiovascular events, 278-285

Caribbean

HHV-6 antibody titers in, 497-500

Carotenoids

consumption of

in fruits and vegetables, 255-260, 333-335 (editorial)

Carotid artery

intima-media thickness

association with low serum albumin, 186-191

Case-control studies

of bladder cancer, 427-431

of breast cancer, 315-320

of childhood cancer

and parental use of alcohol and tobacco, 354-359

of Hodgkin's disease and Vietnam service, 400–406

of lifetime occupational physical activity and hip fracture risk in women, 321–324

of maternal education correlated with mental retardation

in 10-year-old children, 347-353

of ovarian cancer, 310-314

response rates among control subjects in, 245-249 on selected diseases and cataract risk in women, 234-238

Caucasians

coronary heart disease and stroke-related mortality for in California, 1985 to 1990, 432-439

and correlates of body fat distribution, 192-200

and long-term survival after coronary heart disease, 25-32

mothers

and maternal education correlated with mental retardation in 10-year-old children, 347-353

and nutrient intakes of 9- and 10-year-old girls, 360-368

perinatal mortality rates in, 393-399

thrombolysis in, 171-178, 250-252 (editorial)

women

role of marriage and socioeconomic status on depression in, 455–463

Cerebrovascular disorders. see also Stroke

mortality from

for six ethnic groups in California, 1985 to 1990, 432–439

Cervical cancer

Trichomonas vaginalis in, 325-332

Changes in incidence

in interpreting HIV sentinel seroprevalence studies, 447–454

Changing of the guard (editorial), 82

Chemical ototoxicity

styrene associated with, 15-24

Children

cancer in

and parental use of alcohol and tobacco, 354–359 and HHV-6 antibody titers in West Africa and the Caribbean, 497–500 and nutrient intakes of 9- and 10-year-old girls, 360–

368

7 to 12 years old

food frequency questionnaires for, 369-377

ten-year-old

maternal education correlated with mental retardation in, 347-353

Chinese populations

coronary heart disease and stroke-related mortality for in California, 1985 to 1990, 432-439

Chinese prospective study

of Trichomonas vaginalis in cervical cancer, 325-332

Cholesterol

and food-frequency questionnaire among

Mexican-Americans in Starr County, Texas, 378-385

in Greek adolescents

relation to diet, obesity, and socioeconomic factors, 286-291

vitamin C supplementation effect on, 52-59 52

Chronic diseases

major

HDL cholesterol association with in US and Russian men, 179-185

Clinical trials

DASH trial fror lowering hypertension, 108–118 phase I hypertension prevention trials, 85–95, 96–107 phase II hypertension prevention trials, 140–148, 149–155, 156–164

TOHP phase II (editorial), 165–167 TONE trial, for blood pressure control, 119–129

Coffee consumption

bladder cancer in relationship to 427-431

Cognitive function

decline over one year interval

assessed via epidemiologic catchment study, 1-7, 1-7 (editorial), 76-78

Cohort studies

assessing cardiovascular disease prevalence in elderly, 270-277

comparing standardized mortality ratios, 60-64 of coronary heart disease and stroke in elderly, 278-285 of diverticular disease risk in men, 221-228

HIV sentinel seroprevalence studies based on, 447–454 of long-term survival after coronary heart disease, 25–32

of Trichomonas vaginalis in cervical cancer, 325-332

Colon

cancer

trends in over a half century in Rochester, Minnesota, 210-214 210

cancer risk

and association of reproductive and menstrual characteristics, 303-309

diverticular disease in, 221-228 221

sigmoid adenoma

cigarette smoking, alcohol use, and dietary habits in, 478-483

Commentary-

American College of Epidemiology, statement of principles (ACE section), 503-504

Community practice

and thrombolytic therapy for acute myocardial infarction (editorial), 250-252

Community surveys

of correlates of body fat distribution, 192–200 educational attainment correlated with cognitive function decline, 1–7, 76–78 (editorial)

Comparative studies

of HHV-6 antibody titers

in West Africa and the Caribbean, 497-500 on prevention of hypertension, 149-155

of two methods of lactose consumption measurement, 473-477

Completeness

AIDS-reporting, 337-346

Congestive heart failure

prevalence

assessing in elderly, 270-277

Connective tissue disease

screening questionnaire

for population studies, 297-302

Consumption

lactose

reliability and comparison of two methods, 473-477

Coronary heart disease

in Greek adolescents, 286-291

HDL cholesterol association with

long term survival after, 25-32

in US and Russian men, 179-185

mortality

for six ethnic groups in California, 1985 to 1990, 432–439

stroke-related mortality

for six ethnic groups in California, 432-439

surveillance and ascertainment of

in Cardiovascular Health Study, 278-285

in women

complexity of studying (editorial), 79-81

Corpus Christi Heart Project

on use and effectiveness of thrombolytic therapy, 171–178, 250–252 (editorial)

Correlates

of body fat distribution, 192-200

Crude nutrient intake

and reproducibility of Italian food frequency questionnaire, 69

Cystitis

history of

bladder cancer in relationship to 427-431

D

Data collection

for cardiovascular events

in Cardiovascular Health Study, 278-285

Deaths. see also Mortality

comparing standardized mortality ratios, 60-64

in interpreting HIV sentinel seroprevalence studies, 447–454

Depression

in black and white women

role of marriage and socioeconomic status, 455-463

Diabetes mellitus

and cataract risk in women, 234-238

incidence and predictors

in Japanese-American men, 33-43

low serum albumin associated with, 186-191

Diagnostic codes

for AIDS surveillance, 337-346

Diary diet listing lactose-containing foods, 473-477 Diet approaches in DASH program for lowering blood pressure, 108attributable risks for bladder cancer in northern Italy 427-431 diary listing lactose-containing foods, 473-477 food frequency questionnaire designed for use in girls age 7 to 12 years, 369-377 habits and sigmoid colon adenomas, 478-483 Mediterranean relation of blood lipids to in Greek adolescents, 286repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire, 65reproducibility of food frequency questionnaire for cancer studies, 69 validity of 24-hour dietary recall interviews conducted among volunteers in adult working community, 484-489 Dietary recall interviews 24-hour conducted among volunteers in adult working community, 484-489 Dietary Approaches to Stop Hypertension (DASH) trial clarification of term ideal in article on (letter), 502 rationale and design of, 108-118 Digestive tract cancer and reproducibility of Italian food frequency questionnaire, 69 Discharge data hospital for AIDS surveillance, 337-346 Distantly recalled activity reliability of, 292-296 Diverticular disease risk of symptomatic in men, 221-228

E Educational level correlated with decline in cognitive function, assessed via epidemiologic catchment study, 1-7, 76-78 (editorial) maternal correlated with mental retardation in 10-year-old children, 347-353

parental and nutrient intakes of 9- and 10-year-old girls, 360-Elderly assessing cardiovascular disease prevalence in, 270-277 carotenoid consumption by in fruits and vegetables, 255-260, 333-335 (editorial) hypertension in, 44-51 repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire estimated in, 65-68 TONE trial for blood pressure control in, 119-129 Environmental factors for HHV-6 antibody titers in West Africa and the Caribbean, 497-500 Epidemiologic catchment study education and change in cognitive function, 1-7, 76-78 (editorial) Erratum, 418 Erythocyte cations in hypertension, 44-51 Estrogen replacement therapy protects against coronary heart disease in women, 25-Ethanol. see also Alcohol and repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire 65-Etiology of hypertension, 44-51 Exercise. see also Exertion; Physical activity and correlates of body fat distribution, 192-200 and incidence and predictors of diabetes in Japanese-American men, 33-43 Exertion. see also Exercise; Physical activity

F Farm machinery occupational injuries related to, 419-426 Fat, body. see also Obesity correlations of distribution, 192-200 Fat, dietary and food-frequency questionnaire among Mexican-Americans in Starr County, Texas, 378and nutrient intakes of 9- and 10-year-old girls, 360and repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire 65and reproducibility of Italian food frequency questionnaire, 69

and hip fractures in women, 321-324

Follow-up methods

for surveillance and ascertainment of cardiovascular events, 278-285

Food-frequency questionnaires

on blood lipids in Greek adolescents, 286-291

designed for use in girls age 7 to 12 years, 369-377

how far have we come and where are we going? (editorial), 413

and lactose consumption measurement, 473-477

for low-income Mexican-Americans in Starr County, Texas, 378–385

quantitative

reproducibility of for cancer studies, 69

semiquantitative

repeatability of nutrient intakes by eledely subjects estimated by, 65-68

Free-living populations

validity of dietary data collected from, 484-489

Frequency-only questions

and reproducibility of Italian food frequency questionnaire, 69

Fruits

carotenoid consumption in and decreased cardiovascular mortality in elderly, 255–260, 333–335 (editorial)

G

Gallbladder disease

and food-frequency questionnaire among

Mexican-Americans in Starr County, Texas, 378-

nonneoplastic, trends in mortality from, 215–220 Generalizibility

in interpreting HIV sentinel seroprevalence studies, 447-454

Geometric mean titer

HHV-6 antibody

in West Africa and the Caribbean, 497-500

Gestational age

and disparity between Australian Aboriginal and Caucasian perinatal mortality rates, 393-399

Ghana

HHV-6 antibody titers in, 497-500

Greek adolescents

blood lipids in

relation to diet, obesity, and socioeconomic factors, 286-291

Growth factors

uterine

in ovarian cancer pathogenesis, 310-314

Gubbio Population Study

of effect of red blood cell sodium and potassium concentration on blood pressure, 44-51

H

Habits

dietary

resulting in sigmoid colon adenomas, 478-483

HDL. see High-density lipoprotein

Health

occupational

and styrene-associated ototoxicity, 15-24

Hearing loss

bone mass values related to

in elderly rural women, 8-14

styrene exposure associated with, 15-24

Heart disease

cancer and stroke, total mortality and mortality from, from 1950 to 1987 in 27 countries (book review), 417

HHV-6. see Human herpesvirus-6

High-density lipoprotein (HDL) cholesterol

in Greek adolescents

relation to diet, obesity, and socioeconomic factors, 286-291

in major noncommunicable diseases

in samples of US and Russian men, 179-185

vitamin C supplementation effect on, 52-59

zinc supplementation associated with higher levels of, 490-496

Hip fractures

in women

risk of, 321-324

Hispanics. see also Mexican Americans

coronary heart disease and stroke-related mortality for in California, 1985 to 1990, 432-439

thrombolysis in, 171-178, 250-252 (editorial)

HIV. see Human immunodeficiency virus

Hodgkin's disease

correlated with Vietnam service, 400-406

Honolulu Heart Program

on incidence and predictors of diabetes in

Japanese-American men, 33-43

Hormone replacement therapy

and thyroid antibody prevalence

in middle-aged women, 229-233

Hospital discharge data

adminstratively collected

for AIDS surveillance, 337-346

Host factors

for HHV-6 antibody titers in West Africa and the Caribbean, 497-500

Household income

and nutrient intakes of 9- and 10-year-old girls, 360-368

Human herpesvirus-6 (HHV-6)

antibody titers

in West Africa and the Carribean, 497-500

Human immunodeficiency virus (HIV) AIDS surveillance administratively collected hospital discharge data for, 337-346 prevalence estimating from sentinel seroprevalance studies, 447-454 Hyperlipidemia and cataract risk in women, 234-238 234 Hypertension. see also Blood pressure and cataract risk in women, 234-238 DASH trial for lowering, 108-118 red blood cell sodium and potassium concentration in 44-51 TONE trial for lowering, 119–129 24-hour dietary recall interviews conducted among volunteers in adult working community, 484-489 Hypertension prevention trials phase II baseline characteristics of participants in, 149-155 calcium and magnesium supplementation in adults with high-normal blood pressure, 86-107 design of, 140-148 potassium supplementation in persons with high-normal blood pressure, 85-95 recruitment for, 140-148 weight loss and dietary sodium reduction interventions, 156-164 Hypnotics prescription for risk of traffic accident injury after, 239-244 Hypothetical cohorts HIV sentinel seroprevalence studies based on, 447-454 Hysterectomy prior associated with ovarian cancer, 310-314 I

prescription for
risk of traffic accident injury after, 239–244

Hypothetical cohorts
HIV sentinel seroprevalence studies based on, 447–454

Hysterectomy
prior
associated with ovarian cancer, 310–314

I
Ideal
clarification of as term
in rationale and design of DASH trial (letter), 502

IFA. see Immunofluorescence assay

Immunofluorescence assay (IFA)
of HHV-6 antibody titers in West Africa and the
Caribbean, 497–500

Inaccuracy
of self-reported weights and heights among American
Indian adolescents, 386–392

Incidence
of colorectal cancer
over a half century in Rochester, Minnesota, 210–
214

of diabetes mellitus in Japanese-American men, 33-43 Incompletely defined cohorts HIV sentinel seroprevalence studies based on, 447-454 Injury Severity Ccore (ISS) importance and uses, 440-446 440 Internal validity in interpreting HIV sentinel seroprevalence studies, International Classification of Diseases for AIDS surveillance, 337-346 Intima-media thickness carotid artery association with low serum albumin, 186-191 Intravenous drug users AIDS in, 337-346 ISS. see Injury Severity Score Italy attributable risks for bladder cancer in 427-431 red blood cells and potassium concentration effects on blood pressure in, 44-51 reproducibility of food frequency questionnaire for cancer studies, 69 selected diseases and cataract risk in women, 234-238

Jamaica
HHV-6 antibody titers in, 497–500
Japan
sigmoid colon adenomas in, 478–483
Japanese-Americans. see also Asian-Americans
coronary heart disease and stroke-related mortality for
in California, 1985 to 1990, 432–439
incidence and predictors of diabetes in, 33–43
Joint classifications
in food frequency questionnaire designed for use in girls
age 7 to 12 years, 369–377

Laboratory problems
in interpreting HIV sentinel seroprevalence studies,
447-454

Lactose consumption
measurement, reliability and comparison of two
methods, 473-477

LDL. see Low-density lipoprotein

Life-style intervention
for prevention and treatment of high blood pressure
(editorial), 165-167

Lifetime occupational physical activity
and hip fracture risk in women, 321-324

Linoleic acid
and reproducibility of Italian food frequency
questionnaire, 69

Lipid(s)

blood, in Greek adolescents, 286-291

body, and cataract risk in women, 234-238

serum profiles in 778 adults, 490-496

Lipoprotein cholesterol

vitamin C supplementation effect on, 52-59

Logic

of modern medicine (book review), 501

Logistic regression model

of educational and cognitive function decline, 1-7

of machine-related occupational injuries in farm

residents, 419-426

Longitudinal studies educational attenment correlated with cognitive

function decline, 1-7, 76-78 (editorial)

Long-term survival

after coronary heart disease, 25-32

Low-density (LDL) lipoprotein cholesterol

in Greek adolescents, relation to diet, obesity, and

socioeconomic factors, 286-291

vitamin C supplementation effect on, 52-59

zinc supplementation associated with higher levels of, 490-496

M

Machine-related injuries

occupational

in farm residents, 419-426

Macular degeneration

age-related

zinc supplementation for, 490-496

Magnesium

supplementation

in persons with high-normal blood pressure, 96-107

Marriage

role of

in depression in black and white women, 455-463

Maternal education

correlated with mental retardation

in 10-year-old children, 347-353

Mediterranean diet

relation of blood lipids to in Greek adolescents, 286-291

Menopause

and thyroid antibody prevalence in middle-aged women, 229-233

Menstruation

and reproductive characteristics

and colon and rectal cancer risk, 303-309

and thyroid antibody prevalence

in middle-aged women, 229-233

Mental retardation

correlated with maternal education

in 10-year-old children, 347-353

Meta-analysis methods

formal

in interpreting HIV sentinel seroprevalence studies,

447-454

Mexican-Americans. see also Hispanics

in Starr County, Texas

food-frequency questionnaire validation among, 378-

385

thrombolysis in, 171-178, 250-252 (editorial)

Migration

in interpreting HIV sentinel seroprevalence studies,

447-454

Mini-Mental State Examination (MMSE)

cognitive function assessed by, in epidemiologic

catchment study, 1-7

Misclassification

in surveillance and ascertainment of cardiovascular

events, 278-285

MMSE. see Mini-Mental State Examination

Modern medicine

logic of, (book review), 501

Morbidity

Injury Severity Score method for retrospective

comparison of, 440-446

Mortality. see also Deaths

cardiovascular, decreased by carotenoid consumption

in fruits and vegetables, 255-260, 333-335 (editorial)

comparing standardized mortality ratios, 60-64

from coronary heart disease and stroke for six ethnic groups in California, 1985 to 1990, 432-439

HDL cholesterol association with

for chronic diseases in US and Russian men, 179-

185

from heart disease, cancer and stroke, from 1950 to 1987

in 27 countries (book review), 417

in interpreting HIV sentinel seroprevalence studies, 447–454

from nonneoplastic gallbladder disease, 215-220

Vietnam service correlated with trauma and selected cancers, 407-412

Mucinous tumors

ovarian

uterine growth factors in pathogenesis of, 310-314

Multicenter controlled feeding study

of dietary patterns to lower blood pressure, 108–118, 130–139

Myocardial infarction

in elderly

decreased by carotenoid consumption in fruits and

vegetables, 255-260, 333-335 (editorial)

long term survival after

comparisons between men and women, 25-32

prevalence assessing in elderly, 270-277

and thrombolytic therapy, 171-178, 250-252 (editorial) complexity of studying (editorial), 79-81

N

Neurologic conditions and maternal education correlated with mental retardation in 10-year-old children, 347-353

NHLBI Growth and Health Study and factors bearing on nutrient intakes of 9- and 10-year old girls, 360-368

Noise exposure

and styrene-associated ototoxicity, 15-24

Noncommunicable diseases

major

HDL cholesterol association with in US and Russian men, 179-185

Nonpharmacological intervention

for blood pressure control in elderly, 119-129 for hypertension prevention, 156-164

Nonprobability samples

HIV sentinel seroprevalence studies based on, 447-454

Nutrient values

estimated from semiquantitative food frequency questionnaire repeatability of, 65-68

O

Obesity. see also Body weight, Fat, body in American Indian adolescents, inaccuracy in selfreporting of, 386-392 386 and cataract risk in women, 234-238 and incidence and predictors of diabetes in Japanese-American men, 33-43 relation of blood lipids to in Greek adolescents, 286-291

Occupational exposures

bladder cancer in relationship to 427-431 and styrene-associated ototoxicity, 15-24

Occupational injuries

machine-related in farm residents, 419-426

Older adults. see Elderly

Oleic acid

and reproducibility of Italian food frequency questionnaire, 69

One-year interval

educational attainment and cognitive function decline over, assessed via epidemiologic catchment study, 1-7, 76-78 (editorial)

Open questions

and reproducibility of Italian food frequency questionnaire, 69

Osteoporosis

and hip fractures in women, 321-324

Ototoxicity

chemical

styrene associated with, 15-24

Ovarian cancer

pathogenesis

epidemiologic evidence for uterine growth factors in, 310-314

and reproducibility of Italian food frequency questionnaire, 69

Overall injury data

Injury Severity Score method for retrospective comparison of, 440-446

P

Paired t test

and repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire 65-

Parental education

and nutrient intakes of 9- and 10-year-old girls, 360-368

Pearson correlation coefficients

and reproducibility of Italian food frequency questionnaire, 69

Pelvic surgery

associated with ovarian cancer, 310-314

Perinatal mortality

Australian Aboriginal and Caucasuna, disparity between, 393-399

Pharmacoepidemiology

and risk for traffic accident injury after prescription for benzodiazepine, 239-244

Phenoxy herbicides

as Hodgkin's disease factors, 400-406

Physical activity. see also Exercise; Exertion

lifetime occupational

and hip fracture risk in women, 321-324

reliability of recall of, 292-296

Placebo-controlled trial

vitamin C supplementaion

effect of on lipoprotein cholesterol, apoilipoprotein, and triglycerides, 52-59

Plasma

ascorbic acid effect of on lipoprotein cholesterol, apolipoprotein, and triglycerides, 52-59

Plastics industry

and styrene-associated ototoxicity, 15-24

Poisson regression

for comparing standardized mortality ratios, 60-64

Polyunsaturated fats (PUFAs)

and nutrient intakes of 9- and 10-year-old girls, 360-368

and reproducibility of Italian food frequency questionnaire, 69

Population studies

comparing standardized mortality ratios, 60-64

of coronary heart disease and stroke in elderly, 278-285

of correlates of body fat distribution, 192-200

of use and effectiveness of thrombolytic therapy, 171–178, 250–252 (editorial)

Postmenopause

and long-term survival after coronary heart disease, 25–32

Potassium

and nutrient intakes of 9- and 10-year-old girls, 360-368

in red blood cells, effect of on blood pressure, 44-51 supplementation

in persons with high-normal blood pressure, 85-95

Predictors

diabetes mellitus

in Japanese-American men, 33-43

of randomization

for phase II hypertension preention trials, 140

Prematurity

and disparity between Australian Aboriginal and Caucasian perinatal mortality rates, 393-399

Presbycusis

in elderly rural women, 8-14

Prevalence

of cardiovascular disease

assessing in elderly, 270-277

in HIV sentinel serosurveys, 447-454

of thyroid antibodies

among healthy middle-aged women, 229-233

Prevention

of hypertension

trials of, 85-95, 96-107, 130-139, 140-148, 149-155, 156-164,

Proportionality assumption

in comparing standardized mortality ratios, 60-64

Prospective studies

of consumption of carotenoids in fruits and vegetables, and decreased cardiovascular mortality in elderly, 255– 260, 333–335 (editorial)

of diverticular disease risk in men, 221-228

of Trichomonas vaginalis in cervical cancer, 325-332

Psychoactive drugs

prescription for

risk of traffic accident injury after, 239-244

Public school system records

and maternal education correlated with mental retardation in 10-year-old children, 347-353

PUFAs. see Polyunsaturated fatty acids

O

Quadratic k values

and repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire 65–68

Quantitative food frequency questionnaire reproducibility of for cancer studies, 69

Questionnaires

food-frequency

on blood lipids in Greek adolescents, 286-291 designed for use in girls age 7 to 12 years, 369-377 how far have we come and where are we going? (editorial), 423

and lactose consumption measurement, 473–477 validation among Mexican-Americans in Starr County, Texas, 378–385

quantitative food frequency

reproducibility of for cancer studies, 69

screening

for connective tissue disease, 297-302 semiquantitative food frequency

nutrition values estimated from, 65-68

Quintile categories

and repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire 65-

R

Race factors

and correlates of body fat distribution, 192-200 and depression in women, 455-463

and maternal education correlated with mental retardation in 10-year-old children, 347-353

and nutrient intakes of 9- and 10-year-old girls, 360-368

Randomization strategies

for phase II hypertension prevention trials, 140-148 Randomized trials

of antioxidants and cardiovascular disease in women, 261-269

applying lessons from to community practice and thrombolytic therapy for acute myocardial infarction (editorial), 250–252

Rate ratio

Vietnam service and risk of death from trauma and selected cancers, 407-412

Ratios

standardized mortality, 60-64

Recall interviews 24-hour dietary, 484-489 of physical activity of several years ago reliability of, 292-296 Recruitment for phase II hypertension prevention trials, 140-148 Rectal cancer risk and association of reproductive and menstrual characteristics, 303-309 Red blood cells potassium and sodium in effect of on blood pressure, 44-51 Reference population in comparing standardized mortality ratios, 60-64 Rehabilitation Injury Severity Score method for retrospective comparison of, 440-446 Reliability of food-frequency questionnaire among Mexican-Americans in Starr County, Texas, 378in interpreting HIV sentinel seroprevalence studies, 447-454 of lactose consumption measurement, 473-477 of methods of assessing cardiovascular disease prevalence in elderly, 270-277 of recall of physical activity, 292-296 Repeatability of food frequency questionaire designed for use in girls age 7 to 12 years, 369-377 of nutrient values estimated from semiquantitative food frequency questionnaire, 65-68 Reproductive factors and breast cancer risk effect of age at diagnosis, 315-320 and menstrual characteristics and colon and rectal cancer risk, 303-309 Response rates among control subjects in case-control studies, 245-249 Retrospective cohort comparison on terfenadine-associated ventricular arrhythmias QTc interval prolongation with, 201-209 Retrospective comparison of overall injury data Injury Severity Score method for, 440-446 Retrospective recall of physical activity of several years ago, 292-296 Risk factors

for bladder cancer in northern Italy, 427-431

and reproductive factors, 315-320

breast cancer

in women, 234-238

cataract

colon and rectal cancer association of reproductive and menstrual characteristics in, 303-309 for diverticular disease in men, 221-228 for HDL cholesterol association with chronic diseases in US and Russian men, 179-185 hip fractures in women, 321-324 Hodgkin's disease and Vietnam service, 400-406 and incidence and predictors of diabetes in Japanese-American men, 33-43 and long-term survival after coronary heart disease, 25-32 and machine-related farm injury, 419-426 repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire, 65for traffic accident injury after prescription for benzodiazepine, 239-244 Trichomonas vaginalis in cervical cancer, 325-332 Russia HDL cholesterol association with chronic diseases in in men, 179-185 S Sampling and response rates among control subjects in case-control studies, 245-249 Saturated fat and food-frequency questionnaire among Mexican-Americans in Starr County, Texas, 378and nutrient intakes of 9- and 10-year-old girls, 360and repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire 65-Screening questionnaire connective tissue disease for population studies, 297-302 Secondary prevention trial of antioxidants and cardiovascular disease in women, 261-269 Self-reporting of weights and heights among American Indian adolescents, 386-392 Self-selection bias in interpreting HIV sentinel seroprevalence studies, Semiquantitative food frequency questionnaire in elderly repeatability of nutrient intakes estimated by, 65-68

Sentinel seroprevalence studies estimating HIV prevalence from, 447–454 Serum

albumin

associated with diabetes mellitus and other cardiovascular risk factors, 186-191 zinc and lipid profiles in 778 adults, 490-496

Sex factors

and correlates of body fat distribution, 192-200 and long-term survival after coronary heart disease, 25-32

Sigmoid colon adenoma

cigarette smoking, alcohol use, and dietary habits in, 478-483

Smoking

bladder cancer in relationship to 427-431 and correlates of body fat distribution, 192-200 in diverticular disease, in men, 221-228 parental

correlated with childhood cancer, 354-359 sigmoid colon adenoma relationship to, 478-483

Social support

as factor relating to depression in Black and White women, 455-463

Sociodemographics

repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire, 65–68

Socioeconomic factors

and nutrient intakes of 9- and 10-year-old girls, 360-368

relation of blood lipids to in Greek adolescents, 286-291

role of

in depression in black and white women, 455-463 Sodium

dietary

reduction of for hypertension prevention, 119-129, 130-139, 140-148, 149-155, 156-164

in red blood cells, effect of on blood pressure, 44-51 Special education services

and maternal education correlated with mental retardation in 10-year-old children, 347-353

Standard error

in interpreting HIV sentinel seroprevalence studies, 447–454

Standardized mortality ratio comparing, 60-64

Starch

in food frequency questionnaire designed for use in girls age 7 to 12 years, 369-377

Statement of principles

Epidemiology and minority populations (ACE section), 505–508

Statistical analysis

comparing standardized mortality ratios, 60-64 and Injury Severity Score, 440-446

Stroke. see also Cerebrovascular disorders

cancer, and heart disease, total mortality and mortality from, from 1950 to 1987 in 27 countries (book review), 417

mortality from for six ethnic groups in California, 1985 to 1990, 432-439

prevalence

assessing in elderly, 270-277

and repeatability of nutrient intakes estimated by semiquantitative food frequency questionnaire, 65– 68

surveillance and ascertainment of in Cardiovascular Health Study, 278–285

Styrene

in hearing loss, 15-24

Subcohorts

comparing standardized mortality ratios, 60-64

Supplementation

calcium

in persons with high-normal blood pressure, 96-107 magnesium

in persons with high-normal blood pressure, 96–107 potassium

in persons with high-normal blood pressure, 85-95 vitamin C

effect of on lipoprotein cholesterol, apolipoprotein, and triglycerides, 52–59

zinc

can adversely affect serum lipid profiles 490–496 Surveillance

AIDS

administratively collected hospital discharge data for, 337–346

of cardiovascular events

in Cardiovascular Health Study, 278-285

Survey methods

in interpreting HIV sentinel seroprevalence studies, 447-454

Survival

Injury Severity Score method for retrospective comparison of, 440-446

long-term

after coronary heart disease, 25-32

Syncope

and terfenadine-associated ventricular arrhythmias, QT_c interval prolongation with, 201–209

T

Terfenadine-associated ventricular arrhythmias and QT_c interval prolongation, 201–209

Triglycerides

vitamin C supplementation effect on, 52-59

Thrombolytic therapy
population-based assessment of use and effectiveness of,
171-178, 250-252 (editorial)
Thyroid antibody prevalence
among healthy middle-aged women, 229-233
Time-averaging
of prevalence in HIV sentinel serosurveys, 447–454
Time or space
populations differing in
Injury Severity Score as method for retrospective
comparison of overall injury data between, 440-
Time trends
in mortality from nonneoplastic gallbladder disease, 215–220
Tobacco use
parental
correlated with childhood cancer, 354–359
TOHP. see Trials of Hypertension Prevention
TONE. see Trial of Nonpharmacological Intervention in
the Elderly
Total mortality
from heart disease, cancer and stroke, from 1950 to 1987
in 27 countries (book review), 417
Traffic accident injury risks of
after prescription for benzodiazepine, 239–244
Transient ischemic attack
prevalence
assessing in elderly, 270–277
Traumatic death
Vietnam service and risk of death from, 407–412
Trends
in colorectal cancer
over a half century in Rochester, Minnesota, 210-
214
in mortality from nonneoplastic gallbladder disease, 215-220
Trial of Nonpharmacological Intervention in the Elderly
(TONE)
design and rationale of blood pressure control trial, 119-
129
Trials of Hypertension Prevention (TOHP)
phase II
baseline characteristics of participants in, 149-155
design of, 140-148
recruitment for, 140-148
weight loss and dietary sodium reduction
interventions, 156–164
Trichomonas vaginalis
in cervical cancer, 325-332

zinc supplementation associated with higher levels of, 490-496 Trinidad HHV-6 antibody titers in, 497-500 Tubal ligation prior associated with ovarian cancer, 310-314 Twenty four-hour dietary recall interviews validity of conducted among volunteers in adult working community, 484-489 U Unbiased estimate comparing standardized mortality ratios, 60-64 United States HDL cholesterol association with chronic diseases in in men, 179-185 Univariate analyis of effect of red blood cell sodium and potassium concentration on blood pressure, 44-51 Urinary tract infection history of bladder cancer in relationship to 427-431 Uterine growth factors epidemiologic evidence for in ovarian cancer pathogenesis, 310-314 Validity of food frequency questionaires among Mexican-Americans in Starr County, Texas, designed for use in girls age 7 to 12 years, 369-377 internal in interpreting HIV sentinel seroprevalence studies, 447-454 of methods of assessing cardiovascular disease prevalence in elderly, 270-277 of 24-hour dietary recall interviews conducted among volunteers in adult working community, 484-489 Vegetable fat and reproducibility of Italian food frequency questionnaire, 69 Vegetables carotenoid consumption in and decreased cardiovascular mortality in elderly, 255-260, 333-335 (editorial) low intake of bladder cancer in relationship to 427-431 Ventricular arrhythmias terfenadine-associated

and QT_c interval prolongation, 201-209

Vietnam service epidemiologic research on difficulties and lessons learned (editorial), 414-416 Hodgkin's disease related to, 400-406 trauma and selected cancers correlated with risk of death from, 407-412 Vitamin B1 in food frequency questionnaire designed for use in girls age 7 to 12 years, 369-377 Vitamin B2 in food frequency questionnaire designed for use in girls age 7 to 12 years, 369-377 Vitamin C. see also Ascorbic acid for cardiovascular disease prevention in women, 261in food frequency questionnaire designed for use in girls age 7 to 12 years, 369-377 and nutrient intakes of 9- and 10-year-old girls, 360supplementation effect of on lipoprotein cholesterol, apolipoprotein, and triglycerides, 52-59 Vitamin E for cardiovascular disease prevention in women, 261and reproducibility of Italian food frequency questionnaire, 69 Vitamins antioxidant for cardiovascular disease prevention in women, 261-269 Volunteers for 24-hour dietary recall interviews, 484-489 Vulnerability to bias comparing standardized mortality ratios, 60-64 in interpreting HIV sentinel seroprevalence studies, 447-454

W

Weight loss

for hypertension prevention, 119-129, 130-139, 140-148, 149-155, 156-164

West Africa

HHV-6 antibody titers in, 497-500

Whites. see Caucasians

Within-stratum-ratio

and standardized mortality ratios, 60-64

Women, Women's health

AIDS in, 337-346

black and white

role of marriage and socioeconomic status on depression in, 455-463

bladder cancer in 427-431 and correlates of body fat distribution, 192-200 Chinese

Trichomonas vaginalis in cervical cancer in, 325-332 coronary heart disease in

complexity of studying (editorial), 79-81

and stroke-related mortality for in California, 1985 to 1990, 432-439

elderly

assessing cardiovascular disease prevalence in, 270-277

elderly rural

age-related hearing losss and bone mass in, 8-14 healthy middle-aged

thyroid antibody prevalence in, 229-233

HHV-6 antibody titers in West Africa and the Caribbean, 497-500

hypertension in, 44-51

hypertension prevention in, 96-107, 149-155

lactose consumption measurement in, 473-477

lifetime occupational physical activity and hip fracture risk in, 321-324

and nutrient intakes of 9- and 10-year-old girls, 360-368

postmenopausal

and long-term survival after coronary heart disease, 25-32

reproductive factors and breast cancer risk in, 315-320 selected diseases and cataract risk in, 234-238

thrombolysis in, 171–178, 250–252 (editorial)

24-hour dietary recall interviews conducted among volunteers in adult working community, 484–489

uterine growth factors in ovarian cancer pathogenesis in, 310-314

vitamin C supplementation in, 52-59

Wisconsin

association of reproductive and menstrual characteristics and colon and rectal cancer risk in, 303-309

Working community

adult

validity of 24-hour dietary recall interviews conducted among volunteers in, 484–489

Z

Zinc

and nutrient intakes of 9- and 10-year-old girls, 360-368

supplements

can adversely affect serum lipid profiles 490-496